

Comparative Study of the Chemical Composition of Three Mango Stem Bark

Victoria Bennett^{1,*}, Azibaola Kesiye Inengite²

¹Department of Chemical Sciences, Faculty of Basic and Applied Sciences, University of Africa, Toru-Orua, Sagbama, Nigeria

²Department of Chemical Sciences, Faculty of Science, Niger Delta University, Amassoma, Nigeria

Email address:

vkalapoi@gmail.com (V. Bennett), Victoria.Bennett@uat.edu.ng (V. Bennett), ainengit@yahoo.com (A. K. Inengite)

*Corresponding author

To cite this article:

Victoria Bennett, Azibaola Kesiye Inengite. Comparative Study of the Chemical Composition of Three Mango Stem Bark. *Journal of Diseases and Medicinal Plants*. Vol. 8, No. 3, 2022, pp. 55-60. doi: 10.11648/j.jdmp.20220803.12

Received: June 12, 2022; Accepted: July 7, 2022; Published: July 20, 2022

Abstract: Mango (*Mangifera indica*) tree is an evergreen domestic plant with immense nutritional and medicinal properties. African mango leaf and root extracts have been reported to have inhibitory activity against several bacteria and fungi. Mango has different species but, this study was focused on the bark of three species (*Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica*). Samples were collected from Toru-Orua and Ebedebiri Communities in Sagbama Local Government Area of Bayelsa State, Nigeria, and a comparative study of the proximate, mineral and bioactive compounds in the bark was carried out. Proximate analysis of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* showed 59.73%, 65.07%, 64.44% moisture content, 1.98%, 4.09%, 2.03% ash content, 1.93%, 1.30%, 1.73% protein content, 5.56%, 6.09%, 7.12% fat content, 0.90%, 0.74%, 2.50% fibre content, and 29.90%, 22.71%, 22.19% carbohydrate content respectively. Result shows that the bark of *Irvingia gabonensis* contains more moisture, hence will be more vulnerable to microbial contamination and high degree of perishability. The bark of *Mangifera indica* contains about three times fiber content than the bark of *Mangifera indica* L. and *Irvingia gabonensis*; implying that the bark of *Mangifera indica* could be a good source of fiber. The value observed for *Irvingia gabonensis* was insignificantly low compared to the fibre content in the seed as reported in the literature. Mineral concentrations of 0.3304%, 0.2296%, 0.2950% K, 0.0056%, 0.0024%, 0.0080% Fe, 0.0050%, 0.0004%, 0.0005% Mn, 1.2163%, 1.1659%, 1.0151% Ca, 0.1684%, 0.2417%, 0.1785% P were analysed respectively. Gas chromatography-mass spectrometry (GC-MS) analysis of methanolic extract of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* showed fifteen, nine and ten bioactive compounds respectively. GC-MS results of the three mango bark species indicate that bioactive compounds are peculiar to a species. Performic acid has been shown to have strong bactericidal and sporicidal effect. Result shows that a significant amount of performic acid, 89.8% is contained in the bark of *Mangifera indica* L. affirming its use in folk medicine. All the analysed compounds have different biological activities justifying the use of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* bark in traditional medicine for the treatment of malaria, typhoid and COVID-19.

Keywords: Mango, Proximate, Mineral, Phytochemicals

1. Introduction

The extensive survey of literature revealed that *Mangifera indica* is an important source of many pharmacologically and medicinally chemicals such as mangiferin, mangiferolic acid, hydroxy-mangiferin, polyphenols and carotenes. Many different pharmacological activities, antioxidant, radio protective, immunomodulatory, anti-allergic, anti-

inflammatory, antitumor, anti-diabetic, lipolytic, antibone resorption, monoamine oxidase-inhibiting, antimicrobial and anti-parasitic, have been reported for mangiferin [1]. All parts are used to treat abscesses, jackal bite, tumor, snakebite, stings, datura poisoning, heat stroke, miscarriage, anthrax, blisters, wounds in the mouth, tympanitis, colic, diarrhea, glossitis, indigestion, bacillosis, bloody dysentery, liver disorders, excessive urination, tetanus and asthma. Mango fruit, seed, leaves, bark and roots have been studied to have

diverse phytochemical constituents including polyphenolic antioxidants, flavonoids, triterpenoids, and micronutrients. Flavor of mango fruits contain volatile organic molecules including terpenes, furanones, lactones and esters. Mango leaves contains multiple minerals and vitamins [2]. African mango leaf and root extracts have documented inhibitory activity against several bacteria and fungi [3, 4]. Potential mechanisms of action include membrane disruption by terpenoids and inactivation of microbial adhesion, enzymes, and cell envelope transport proteins by ellagic acid-like compounds [4].

Rats were fed a normal diet and 1 mL of African mango oil or water over 4 weeks. Abdominal fat was lower, plasma HDL cholesterol and triglyceride levels were higher, and LDL: HDL and total cholesterol: HDL ratios were lower in rats administered the oil.

Blood glucose levels were also lower in rats administered the oil [5]. *Irvingia gabonensis* (IG) seeds and seed extracts have been shown to have hypoglycemic properties as well as antihyperglycemic properties. [6]. The mechanism appears to be associated with (1) down-regulated expression of adipogenic transcription factors or PPAR- γ and adipocyte-specific proteins, such as leptin, and (2) up-regulated expression of adiponectin. Adiponectin has anti-atherogenic, anti-inflammatory, and antidiabetic activity.

A 50% ethanolic extract of the leaves produced a significant hypoglycemic effect at a dose of 250 mg/kg, both in normal and streptozotocin-induced diabetic animals [7]. The stimulation of β -cells to release insulin was thought to be part of the mechanism of action. The effect of the aqueous extract of the leaves on blood glucose level in normoglycaemic, glucose induced hyperglycemic and streptozotocin (STZ)-induced diabetic rats has been assessed. The results indicate that the aqueous extract of the leaves of MI possess hypoglycemic activity. The stem-bark of aqueous extract was used to examine the anti-inflammatory, analgesic and anti-diabetic properties. The different chemical constituents of the plant, especially the polyphenolics, flavonoids, triterpenoids, mangiferin, and other chemical compounds present in the plant may be involved in the observed anti-inflammatory, analgesic, and hypoglycemic effects of the plant's extract [7]. Diabetic animals eating diets containing 5, 10 and 15% mango flour showed significant decrease in blood glucose level [8].

African mango reduces body weight, body fat, waist circumference, blood sugar, triglycerides, and cholesterol [9]. Patients received either 150 mg of African mango seed extract or placebo 30 minutes before lunch and dinner. Patients receiving the extract improved both weight reduction (body weight, body fat, waist circumference) and metabolic parameters (plasma total cholesterol, LDL cholesterol, blood glucose, and C - reactive protein, adiponectin, and leptin levels). Numerous studies exist on the potential industrial application of African mango in food, cosmetic, and pharmaceutical products [10-16]. The seeds are ground into a paste, also known as dika bread, which is valued for its food-thickening properties [17].

Margarine-based African mango oil may provide an alternative to trans-fatty acids obtained during hydrogenation used in oil technological applications [18]. Oral administration of an African mango methanol extract at a dose of 150 and 250 mg/kg significantly ($P < 0.0021$) lowered plasma glucose levels in diabetic rats within 2 hours after treatment. The mechanism of action may involve extract stimulation of pancreatic beta-cell function or hypoglycemic activity via an extra-pancreatic mechanism [9].

Although, manifold studies on the phytochemical composition of different Mango species have been reported in the literature, a comparative study of the biochemical components of the bark of different mango species is yet to be reported.

2. Materials and Methods

2.1. Materials

All chemicals used were of analytical grades and obtained from BDH, Labtech chemicals and used without further purification.

2.2. Methods

Samples of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* bark were collected from Toru-Orua and Ebedebiri Communities in Sagbama Local Government Area of Bayelsa State, Nigeria and properly identified at the Biological Sciences Department of the University of Africa, Toru-Orua. *Mangifera indica* L bark were air dried for fourteen days and pulverized manually and electronically, and stored in a desiccator for analysis.

2.2.1. Proximate Analysis

Standard procedures as described by the Association of Official Analytical Chemists [19] were used in the determination of moisture, Fat, Ash, Crude fibre, Crude Protein and Carbohydrate Content.

2.2.2. Mineral Analysis

Standard procedures as described by the Association of Official Analytical Chemists [19] were used to determine Manganese (Mn), Iron (Fe), Calcium (Ca), Phosphorus (P) and Potassium (K).

2.2.3. Bioactive Chemicals

GC-MS analysis of methanol extract of *Mangifera indica* L bark was performed with GC (Agilent 6890) and MS (5973 MSD) equipped with Restek capillary column (30 m \times 0.53mm; film thickness 0.12 μ m), using Helium as the carrier gas with a flow rate of 1 mL/min.

3. Results

Results of the various analyses carried out on the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* are presented as follows:

Table 1. Proximate Analyses.

| Proximate Principles (%) | Mango species | Mean \pm S.D |
|--------------------------|---------------------|-----------------|
| Moisture | Mangifera indica L. | 59.73 \pm 01 |
| | Irvinga gabonensis | 65.07 \pm 01 |
| | Mangifera indica | 64.44 \pm 01 |
| Ash | Mangifera indica L. | 1.98 \pm 01 |
| | Irvinga gabonensis | 4.09 \pm 01 |
| | Mangifera indica | 2.03 \pm 01 |
| Protein | Mangifera indica L. | 1.93 \pm 0.1 |
| | Irvinga gabonensis | 1.30 \pm 0.1 |
| | Mangifera indica | 1.73 \pm 0.1 |
| Fat | Mangifera indica L. | 5.56 \pm 0.1 |
| | Irvinga gabonensis | 6.09 \pm 0.1 |
| | Mangifera indica | 7.12 \pm 0.1 |
| Fibre | Mangifera indica L. | 0.90 \pm 0.1 |
| | Irvinga gabonensis | 0.74 \pm 0.1 |
| | Mangifera indica | 2.50 \pm 0.1 |
| Carbohydrate | Mangifera indica L. | 29.90 \pm 0.1 |
| | Irvinga gabonensis | 22.71 \pm 0.1 |
| | Mangifera indica | 22.19 \pm 0.1 |

Table 2. Mineral analysis of 3 species of mango barks per 100 grams.

| Parameters (%) | Mango Variety | Mean \pm S.D |
|----------------|---------------------|------------------|
| Potassium (K) | Mangifera indica L. | 0.3304 \pm 0.1 |
| | Irvinga gabonensis | 0.2296 \pm 0.2 |
| | Mangifera indica | 0.2950 \pm 0.1 |
| Iron (Fe) | Mangifera indica L. | 0.0056 \pm 0.2 |
| | Irvinga Gabonensis | 0.0024 \pm 0.2 |
| | Mangifera indica | 0.0080 \pm 0.1 |
| Manganese (Mn) | Mangifera indica L. | 0.0050 \pm 0.1 |
| | Irvinga gabonensis | 0.0004 \pm 0.1 |
| | Mangifera indica | 0.0005 \pm 0.1 |
| Calcium (Ca) | Mangifera indica L. | 1.2163 \pm 0.1 |
| | Irvinga gabonensis | 1.1659 \pm 0.2 |
| | Mangifera indica | 1.0151 \pm 0.1 |
| Phosphorus (P) | Mangifera indica L. | 0.1684 \pm 0.2 |
| | Irvinga gabonensis | 0.2417 \pm 0.2 |
| | Mangifera indica | 0.1785 \pm 0.1 |

Table 3. GC-MS analysis of bioactive components in the bark of *Mangifera indica* L. (German/Opororo) powder.

| S/N | Retention time (Min) | Name of Bioactive compound | Molecular formula | Molecular Weight | Peak area Percent (%) |
|-----|----------------------|--|--|------------------|-----------------------|
| 1. | 1.902 | Performic acid | C ₄ H ₁₀ O ₃ Si | 134 | 89.8 |
| 2. | 6.571 | Fluoroacetic acid | C ₆ H ₉ FO ₄ | 164 | 0.89 |
| 3. | 7.000 | Cyclohexane, 1-ethenyl-1-methyl-2-(1-methylethenyl)-4-(1-methylethylidene)- | C ₁₅ H ₂₄ | 204 | 83.4 |
| 4. | 7.251 | Caryophyllene | C ₁₅ H ₂₄ | 204 | 56.1 |
| 5. | 7.314 | Naphthalene, 1,2,3,4,4a,5,6,7-octahydro-4a,8-dimethyl-2-(1-methylethenyl)- | C ₁₅ H ₂₄ | 204 | 67.1 |
| 6. | 7.560 | 1,4,7,-Cycloundecatriene, 1,5,9,9-tetramethyl-, Z,Z,Z | C ₁₅ H ₂₄ | 204 | 11.5 |
| 7. | 8.509 | Humulene | C ₁₅ H ₂₄ | 204 | 12.2 |
| 8. | 8.926 | Naphthalene, 1,2,3,4,4a,5,6,8a-octahydro-4a,8-dimethyl-2-(1-methylethenyl)-, [2R-(2 α ,4 α ,8 α)]- | C ₁₅ H ₂₄ | 204. | 57.0 |
| 9. | 9.280 | Cyclohexanemethanol | C ₇ H ₁₄ O | 114 | 89.0 |
| 10. | 9.480 | Aromandendrene | C ₁₅ H ₂₄ | 204 | 47.4 |
| 11. | 9.538 | 2-Naphthalenemethanol | C ₁₁ H ₁₀ O | 158 | 77.0 |
| 12. | 9.646 | 1-Naphthalenemethanol | C ₁₁ H ₁₀ O | 158 | 20.2 |
| 13. | 9.720 | Valerena-4,7(11)-diene | C ₁₅ H ₂₄ | 204 | 44.8 |
| 14. | 11.178 | α -Guaiene | C ₁₅ H ₂₄ | 204 | 2.86 |
| 15. | 11.824 | (1R,4aR,7R,8aR)-7-(2-Hydroxypropan-2-yl)-1,4a-dimethyldecahydronaphthalen-1-ol | C ₁₅ H ₂₈ O ₂ | 240 | 91.4 |

Table 4. GC-MS analysis of bioactive components in the bark of *Irvingia gabonensis* (African bush mango) powder.

| S/N | Retention time (Min) | Name(s) of Bioactive compound | Molecular formula | Molecular Weight | Peak area Percent (%) |
|-----|----------------------|--|--|------------------|-----------------------|
| 1. | 1.908 | psi.,psi.-Carotene | C ₄₁ H ₅₈ O | 566 | 22.0 |
| 2. | 4.417 | 2,7-Diphenyl-1,6-dioxypyridazino [4,5:2',3'] pyrrole[4',5'-d]pyridazine | C ₂₀ H ₁₃ N ₅ O ₂ | 328 | 19.4 |
| 3. | 6.103 | Exo-norbornanol dimethyl(pentafluorophenyl)silyl ether | C ₁₅ H ₁₇ F ₅ OSi | 426 | 19.7 |
| 4. | 7.560 | α -D-Glucofuranose 6-O-(trimethylsilyl)-, cyclic 1,2:3,5-bis(butylboronate) | C ₁₇ H ₃₄ B ₂ O ₆ Si | 587 | 7.80 |
| 5. | 7.749 | 2-Hexadecanol | C ₁₆ H ₃₄ O | 228 | 14.7 |
| 6. | 9.715 | 1-Hexadecanol-2-methyl | C ₁₇ H ₃₆ O | 154 | 6.67 |
| 7. | 12.132 | β -Acorenol | C ₁₅ H ₂₆ O | 334 | 5.54 |
| 8. | 12.944 | Patchoulene | C ₁₅ H ₂₄ | 734 | 4.47 |
| 9. | 18.196 | 17-Pentatriacontene | C ₃₅ H ₇₀ | 668 | 7.53 |

Table 5. GC-MS analysis of bioactive components in the bark of *Mangifera indica* (Yellow mango) powder.

| S/N | Retention time (Min) | Name of Bioactive compound | Molecular formula | Molecular Weight | Peak area Percent (%) |
|-----|----------------------|---|--|------------------|-----------------------|
| 1. | 0.656 | Benzo[b]fluoranthene | C ₂₀ H ₁₂ | 252 | 62.1 |
| 2. | 1.896 | Methane-d, trichloro | CCl ₃ D | 119 | 99.0 |
| 3. | 3.068 | Benzo[e] pyrene | C ₂₀ H ₁₂ | 252 | 63.5 |
| 4. | 7.114 | Benzo[k]fluoranthene | C ₂₀ H ₁₂ | 252 | 13.0 |
| 5. | 12.944 | 1,2,3-Triazol, 2-(E-4,4-dicyano-3-N-methylanilino-1,3-butadien-1-yl)-4-(methoxycarbonyl)- | C ₁₇ H ₁₄ N ₆ O ₂ | 334 | 98.4 |
| 6. | 13.304 | 2-(E-4,4-Dicyano-1-N-methylanilino-1,3-butadien-1-yl)-4-(methoxycarbonyl)-1,2,3-triazole | C ₁₇ H ₁₄ N ₆ O ₂ | 334 | 0.98 |
| 7. | 14.178 | Cyclodecasiloxane | C ₂₀ H ₆₀ O ₁₀ Si ₁₀ | 740 | 97.7 |
| 8. | 15.544 | Octasiloxane,1,3,3,5,5,7,7,9,9,11,11,13,13,15,15-hexadecamethyl | C ₁₆ H ₅₀ O ₇ Si ₈ | 578 | 0.97 |
| 9. | 16.847 | 28-Nor-17 α (H)-hopane | C ₂₉ H ₅₀ | 262 | 95.9 |
| 10. | 17.407 | 28-Nor-17 β (H)-hopane | C ₂₉ H ₅₀ | 398 | 3.84 |

4. Discussion

Results of the various analyses carried out on the flesh and seeds of *Treculia africana* are presented as follows.

Table 1 shows the primary metabolites in the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica*. The moisture content of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 59.73%, 65.07% and 64.44% respectively. Analysis indicate that the bark of *Irvingia gabonensis* contains more moisture, hence will be more vulnerable to microbial contamination and high degree of perishability [20]. The moisture content of the bark of *Irvingia gabonensis* was lower compared to the ethanol and distilled Water Extract of the cotyledon of *Irvingia gabonensis*, 84.31%, 94.41% [21]. The ash contents of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 1.98%, 4.09% and 2.03% indicating that the bark of *Irvingia gabonensis* contains higher organic matter than the barks of *Mangifera indica* L and *Mangifera indica*. The observed value of ash content in the bark of *Irvingia gabonensis* was significantly higher than in the ethanol extract of the cotyledon, 0.14% [21]. The protein content of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 1.93%, 1.30% and 1.73% respectively. The value for *Irvingia gabonensis* was higher than the value of the ethanol extract of the cotyledon of *Irvingia gabonensis*, 0.14% [21].

The fat content of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 5.56%, 6.09% and 7.12%. This result indicates that the bark of *Mangifera*

indica contains more fat than the barks of *Mangifera indica* L. and *Irvingia gabonensis*. The fiber content of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 0.90%, 0.74% and 2.50%. The bark of *Mangifera indica* contains about three times fiber than the barks of *Mangifera indica* L., *Irvingia gabonensis*; hence, the bark of *Mangifera indica* is a good source of fiber. The value observed for *Irvingia gabonensis* was insignificantly low compared to the amount of fibre in the seed as reported in literature [22]. The carbohydrate content of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 29.90%, 22.71% and 22.19% respectively. Carbohydrate content in the bark of *Irvingia gabonensis* was significantly higher than the values observed in the cotyledon [21].

Table 2 presents the mineral content of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica*. The concentrations of K in the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* 0.3304%, 0.2296% and 0.2950% respectively, indicating higher potassium (K) concentration in the bark of *Mangifera indica* L., than in the bark of *Irvingia gabonensis* and *Mangifera indica*. The concentrations of iron (Fe) in the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 0.0056%, 0.0024% and 0.0080% respectively. Higher concentrations in the pulp of *Mangifera indica* were reported in literature [23]. The concentrations of Mn in the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 0.0050%, 0.0004% and 0.0005% respectively. The concentration of Mn in the bark of *Mangifera indica* L. was lower than the concentration in the

seed kernel [24] The concentrations of Ca in the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 1.2163%, 1.1659% and 1.0151% respectively. The body needs a constant supply of calcium to provide bone strength against compression [25]. The concentration of Ca in the bark of *Mangifera indica* L. was moderately higher than the concentration in the seed kernel [24]. The amount of P in the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* were 0.1684%, 0.2417% and 0.1785% respectively. The bark of *Irvingia gabonensis* has higher P concentration than the bark of *Mangifera indica* L. and *Mangifera indica*. Phosphorus is important to build strong bony structure [26]. The present comparative study confirms the fact that the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* contain minerals that are important for various body functions.

The analysis of the Gas chromatography-Mass spectrometry (GC-MS) of the methanol extracts of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* are presented in Tables 3 – 5. The Bark of *Mangifera indica* L. (German/Opiororo) shows Fifteen phytochemicals with their retention time, molecular weight, molecular formula and peak area percentage, Table 3. The phytochemicals include: Performic acid, 89.8%, Cyclohexane, 1-ethenyl-1-methyl-2-(1-methylethenyl)-4-(1-methylethylidene)-, 83.4%, Caryophyllene, 56.1%, Naphthalene, 1,2,3,4,4a,5,6,7-octahydro-4a,8-dimethyl-2-(1-methylethenyl)-, 67.1%, Cyclohexanemethanol, 89.0%, Aromandendrene, 47.4%, 2-Naphthalenemethanol, 77.0% and (1R,4aR,7R,8aR)-7-(2-Hydroxypropan-2-yl)-1,4a-dimethyldecahydronaphthalen-1-ol, 91.4%. The maximum peak, 91.4% was shown by 2-Naphthalenemethanol, 77.0% and (1R,4aR,7R,8aR)-7-(2-Hydroxypropan-2-yl)-1,4a-dimethyldecahydronaphthalen-1-ol and the lowest peak, 0.89% was shown by Fluoroacetic acid. The bioactive components whose peak area percentages were less than 5% were considered to be insignificant Caryophyllene is one of the chemical compounds that contribute to the sweet smell of the bark of *Mangifera indica* L. Caryophyllene has been given GRAS (generally regarded as safe) designation by the FDA and is approved by the FDA for use as a food additive, typically for flavoring [20].

A significant amount of Performic acid in the bark of *Mangifera indica* L confirms its use in folk medicine. Performic acid has proven strong bactericidal and sporicidal effect [27]. All the different phytochemicals identified from the bark of *Mangifera indica* L were found to possess biological properties such as anti-microbial, anti-cancer, anti-inflammatory and anti-malaria activities which corroborates the use of the bark of *Mangifera indica* L. in folk medicine.

Nine phytochemicals were analysed in the bark of *Irvingia gabonensis* (African bush mango), Table 4. These include: psi.,psi.-Carotene, 22.0%, 2,7-Diphenyl-1,6-dioxypyridazino[4,5:2',3'] pyrrole[4',5'-d]pyridazine, 19.4% and Exo-norbornanol dimethyl(pentafluorophenyl)silyl ether, 19.7%. Maximum peak (22.0%) was shown by psi.,psi.-Carotene and minimum peak (4.47%) by Patchoulene.

Carotenoid pigments present in fruits and vegetables are the main dietary source of Vitamin A [28].

Ten phytochemicals were analysed from the bark of *Mangifera indica* (Yellow mango), Table 5. The maximum peak, 99.0% was shown by Methane-d, trichloro, 99.0% and minimum peak, 0.97% was shown by Octasiloxane, 1,3,3,5,5,7,7,9,9,11,11,13,13,15,15-hexadecamethyl. The GC-MS analyses of the bark of *Mangifera indica* shows bioactive components different from the bioactive components analysed from the bark of *Irvingia gabonensis* and *Mangifera indica* L. These bioactives corroborate their use in folk medicine in curing malaria, typhoid and COVID-19.

5. Conclusion

Comparative study of the Proximate, minerals and bioactive components in methanol extract of the bark of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* was conducted. Results indicate that the bark of *Irvingia gabonensis* contains more moisture, ash and fat than the bark of *Mangifera indica* L. and *Mangifera indica*. Moisture content plays a significant role in the physical appearance, shelf-life and resistance to bacterial contamination of species. Minerals result revealed that the bark of *Irvingia gabonensis* is richer in calcium than the bark of *Mangifera indica* L. and *Mangifera indica*. Gas chromatography-mass spectrometry (GC-MS) analysis of methanolic extract of *Mangifera indica* L., *Irvingia gabonensis* and *Mangifera indica* presented 15, 9 and 10 biochemicals. The bark of *Mangifera indica* L. contains a significant amounts of Performic acid and Caryophyllene. The bark of *Irvingia gabonensis* contains a moderate amount of psi.,psi.-Carotene. Methane-d, trichloro, 1,2,3-Triazol, 2-(E-4,4-dicyano-3-N-methylanilino-1,3-butadien-1-yl)-4-(methoxycarbonyl)-, cyclodecasiloxane and 28-Nor-17 α (H)-hopane. GC-MS results imply that the bark of *Mangifera indica* L, *Irvingia gabonensis* and *Mangifera indica* contain variant bioactive components. All the analysed compounds have different biological activities justifying their use in traditional medicine for the treatment of malaria, typhoid and COVID-19.

6. Recommendation

Extracts of mango stem bark have been used locally to cure COVID-19. This study therefore recommends isolation of the different identified bioactive compounds for medicinal and pharmacological application.

References

- [1] Ronchi S. N., Brasil G. A., Do N., Andrews M., De L., Ewelyne M.(2015) Phytochemical and *in vitro* and *in vivo* biological investigation on the antihypertensive activity of mango leaves (*Mangifera indica* L.) Therapeutic Advances in Cardiovascular Disease, 9 (5), 244-256.

- [2] Kumar M., Saurabh V., Tomar M., Hasan M., Changan S., Sasi M., Maheshwari C., Prajapati U., Singh S., Prajapat R. K., Dhumal S., Punia S., Amarowicz R. and Mekhemar M. (2021) Mango (*Mangifera indica* L.) Leaves: Nutritional Composition, Phytochemical Profile, and Health-Promoting Bioactivities. *Antioxidants* 10 (2), 299; <https://doi.org/10.3390/antiox10020299>
- [3] Fadare D. A. and Ajaiyeoba E. O. (2008) Phytochemical and antimicrobial activities of the wild mango-*Irvingia gabonensis* extracts and fractions. *African Journal of Medicine and Medical Sciences* 37 (2): 119-24.
- [4] Kumar M., Saurabh V., Tomar M. Muzaffar Hasan M. (2021) Mango (*Mangifera indica* L.) Leaves: Nutritional Composition, Phytochemical Profile, and Health-Promoting Bioactivities. *Antioxidants* 10 (2) 299.
- [5] Ogunsina B., Bhatnagar A. S. (2012) Proximate composition of African bush mango (*Irvingia gabonensis*) kernels and characteristics of its oil <https://www.researchgate.net/publication/267336250>
- [6] Oben J. E., (2011) Seed Extract of the West African Bush Mango (*Irvingia Gabonensis*) and its Use in Health. *Nut and seeds in Health and Disease Prevention*, 271-278.
- [7] Shah K. A., Patel M. B., Patel R. J., Parmar P. K. (2010) *Mangifera Indica (Mango)*. *Pharmacognosy Review* 4 (7) 42-48.
- [8] Perpétuo G. F., Salgado J. M. (2003) Effect of mango (*Mangifera indica*, L.) ingestion on blood glucose levels of normal and diabetic rats. *Plant Foods For Human Nutrition* 58, 1-12.
- [9] Rivera V. N., Pérez S. C., Chanona-Pérez J., Perea-Flores M. J., Calderón-Domínguez G., Farrera-Rebollo R (2014) Industrial applications and potential pharmaceutical uses of mango (*Mangifera indica*) kernel In book: *Seeds as Functional Foods and Nutraceuticals: New Frontiers in Food Science*. Chapter: Industrial applications and potential pharmaceutical uses of mango (*Mangifera indica*) kernel.
- [10] Teresa M., Ponce, F., Cardoso L. C., Mantell C., Ossa M. D. (11th Int. Conference on Chemical and Process Technology. At: Milano (Italia), 2013) Potential Use of Mango Leaves Extracts Obtained by High Pressure Extraction Technologies in Cosmetic, Pharmaceuticals and Food Industries DOI: 10.13140/RG.2.1.2736.0883.
- [11] Reguengo L., Barbosa-Pereira L., Rembangouet's W., Bertolino M., Giordano M., Rojo-Poveda O., Zeppa G. (2019) Food applications of *Irvingia gabonensis* (Aubry-Lecomte ex. O'Rorke) Baill., the 'bush mango': A review. *Critical Reviews in Food Science and Nutrition* 60 (3).
- [12] Yoon S., Kim J., Lee B. S., Baek S. C., Chung S. J. and Kim K. H. (2022) Terminalin from African Mango (*Irvingia gabonensis*) Stimulates Glucose Uptake through Inhibition of Protein Tyrosine Phosphatases, *bimolecules* 12, 321.
- [13] Parvez G. M. M. (2016) Pharmacological Activities of Mango (*Mangifera Indica*): A Review. *Journal of Pharmacognosy and Phytochemistry* 5 (3): 01-07.
- [14] Joseph K. and Aworh O. (1992) Post-harvest treatment of wild mango (*Irvingia gabonensis*) for improved shelf life. *Food Chemistry* 44 (1) 45-48.
- [15] Wongkaew M., Chaimongkol P., Leksawasdi N., Jantanasakulwong K., Rachtanapun P., Seesuriyachan P., Phimolsiripol Y., Chaityaso T., Ruksiriwanich W., Jantrawut P. and Sommano S. R. (2021) Mango Peel Pectin: Recovery, Functionality and Sustainable Uses. *Polymers* 13 (22): 3898. doi: 10.3390/polym13223898.
- [16] Umekoli, G. C., Onyechi, J. O. and Udeala, O. K (2009) Use of Dika Fat in the Formulation of Sustained Release Theophylline Tablets and Capsules. *Bio-Research*, 7 (1): 456-460.
- [17] Onyeike E. N and Acheru G. N. (2002) Chemical composition of selected Nigerian oil seeds and physicochemical properties of the oil extracts. *Food Chemistry* 77 (4), 431-437.
- [18] Matos L., Nzikou J. M., Matouba E., Pandzou-Yembe V. N (2009) Studies of *Irvingia gabonensis* Seed Kernels: Oil Technological Applications. *Pakistan Journal of Nutrition* 8 (2).
- [19] AOAC (2005) Official method of Analysis. 18th Ed., Association of Officiating Analytical Chemists, Washington DC, Method 935.14 and 992.24.
- [20] Bennett V., Isaiah T. D. (2022) Proximate, Nutritional and Phytochemical Analysis of *Treculia africana* (African Breadfruit) Decne in South-South, Nigeria. *World Journal of Innovative Research*, 12 (4), 19-23.
- [21] Lawson D., Chioma D. (2018) Proximate Analysis and Phytochemical Screening of *Irvingia Gabonensis* (Agbono Cotyledon). *Biomedical Journal of Scientific and Technical research* 5 (4) DOI 10.26717/BJSTR.2018.05.001227.
- [22] Mgbemena N. M., Ilechukwu I., Okwunodolu F. U., Chukwurah J. O. and Lucky I. B. (2019). Chemical composition, proximate and phytochemical analysis of *Irvingia gabonensis* and *Irvingia wombolu* peels, seed coat, leaves and seeds. *Ovidius University Annals of Chemistry* 30 (1) 65-69.
- [23] Gueye R. S., Ledauphin J., Baraud F., Leleyter L., Lemoine M., Gueye R., Diedhiou A., Balde M., Gaye N., Seck M., Fall D., Wele A. (2022) Evaluation of Mineral Elements Content of Senegal Fruits. *Journal of Mineral and Materials Characterization and Engineering* 10 (2)].
- [24] Chinonye I. I., Chijioke-Okere M., Onu U. L., Ukaoma A. A and Onwumere F. (2018) Minerals and Phytochemical Constituents of *Mangifera indica* Seed Kernel obtained from Eastern Nigeria. *African Journal of Agriculture and Food Science*. 1 (2) 1-11.
- [25] Pu F., Chen N., Xue S. (2016) Calcium intake, calcium homeostasis and health. *Food Science and Human wellness* 5 (1) 8-19.
- [26] Marengo K. LDN (July 1, 2019) What Are the Health Benefits of Phosphorus? *MEDICALNEWTODAY*.
- [27] Kozarov D., Elenkov G., Symnaliev M., Illiev B. (1975) Disinfectant effect of performic acid. *Journal of Hygiene Immunology* 19 (3) 389-92.
- [28] Perera C. O. and Yen G. M. (2007) Functional Properties of Carotenoids in Human Health, *International Journal of Food Properties*, 10:2, 201-230, DOI: 10.1080/10942910601045271.